
Lawn & Garden Pesticides FACTS & FIGURES

PESTICIDE USAGE

- 88 million households in the U.S. use pesticides around their home.¹
- Herbicides account for the highest usage of pesticides in the home and garden sector, with over 28 million pounds applied on lawns and gardens in 2012.²
- Suburban lawns and gardens receive more pesticide applications per acre (3.2-9.8 lbs.) than agriculture (2.7 lbs. per acre on average).³
- Pesticide expenditures (or sales) by the chemical industry average \$9 billion. Annual sales of the landscape industry are over \$53.9 billion.⁴
- Included in the most commonly used pesticides in pounds per year are: 2,4-D (7-9 million), glyphosate/ Roundup (4-6 million), MCPP (Mecoprop) (2-4 million), pendimethalin (2-4 million), carbaryl (2-4 million).⁵
- A 2004 national survey reveals that 5 million homeowners use only organic lawn practices and products and 35 million people use both toxic and non-toxic materials.⁶

HEALTH & EXPOSURE RISKS

Of the 30 commonly used lawn pesticides, 16 are probable or possible carcinogens, 12 are linked with birth defects, 21 with reproductive effects, 14 are neurotoxic, 25 cause liver or kidney damage, 26 are sensitizers and/or irritants, and 17 have the



potential to disrupt the endocrine (hormonal) system.⁷

- Pregnant women, infants and children, the elderly, and the chronically ill are at greatest risk from pesticide exposure, which can increase risk of chronic diseases.⁸
- Scientific studies find significant pesticide residues inside homes due to drift through the air and chemicals tracked in, where they contaminate air, dust, surfaces, and carpets.
 Higher levels of pesticides in a child's home has been associated with higher levels of pesticide residue in their urine.⁹

CHILDREN & PESTICIDES

- Children take in more pesticides relative to body weight than adults and have developing immune, nervous, and digestive systems that make them more vulnerable to environmental toxins.¹⁰
- The President's Cancer Panel on Environmental Cancer Risk notes that leukemia rates are consistently elevated among children whose parents used pesticides in their home and garden.
- The National Academy of Sciences estimates 50% of lifetime pesticide exposure occurs during the first



five years of life.¹¹ A study in Cancer Causes and Control suggests that preconception pesticide exposure and possible exposure during pregnancy is associated with an increased risk of childhood brain tumors.¹²

- Studies show low levels of exposure to lawn pesticide products are linked to increased rates of miscarriage, and suppression of the nervous, endocrine, and immune systems.¹³
- Research finds that young infants and toddlers exposed to herbicides (weedkillers) within their first year of life are 4.5 times more likely to develop asthma by the age of five, and almost 2.5 times more likely when exposed to insecticides.¹⁴
- Researchers at Cincinnati Children's Hospital Medical Center found an association between increasing exposures to commonly used synthetic pyrethroid insecticides and attention deficit/hyperactivity disorder (ADHD), with a higher association in boys than girls.¹⁵
- Children ages 6–11 have higher levels of lawn chemicals in their blood than all other age categories. Biomonitoring studies find that pesticides pass from mother to child through umbilical cord blood and breast milk.¹⁶

WILDLIFE, PETS & PESTICIDES

- A study published in Environmental Research found that dogs whose owners' lawns are professionally treated with pesticides are associated with a significantly higher risk of canine malignant lymphoma.¹⁷
- Of the 30 commonly used lawn pesticides: 22 are toxic to birds, 14 are toxic to mammals, 30 are toxic to fish and aquatic organisms, and 29 are deadly to bees.¹⁸
- Pesticides can be toxic to wildlife and cause food source contamination, behavioral abnormalities that interfere with survival, and death.
- Lawn and garden pesticides are deadly to nontarget species and can harm beneficial insects and soil microorganisms essential to a naturally healthy lawn.²⁰
- Home and garden insecticides in the neonicotinoid class have been linked to pollinator decline, with harm to bees' reproduction mobility, navigation, feeding, foraging, memory and learning.²¹

Lawn and garden pesticides... harm beneficial insects and soil microorganisms essential to a naturally healthy lawn.

PESTICIDES IN THE WATER

- Of the 30 commonly used lawn pesticides, 19 are detected in groundwater, and 20 have the potential to leach.²²
- In a U.S. Geological Survey study of glyphosate contamination in 38 states, scientists detected the chemical in more than 50% of samples of sediment, ditches and drains, precipitation, large rivers, and streams.²³
- Half of shallow wells within the U.S. have detectable levels of pesticides, while 20% of private wells contain at least one contaminant at levels of potential health concern.²⁴
- After the Canadian Province of Ontario implemented restrictions on the use of pesticides, a government report found a staggering decline in herbicide concentrations. Median concentrations for herbicides 2,4-D, dicamba, and MCPP, commonly used by both private individuals and lawn care companies, declined by 81%, 83%, and 71%, respectively.²⁵
- In addition to harming pollinators, neonicotinoid contamination has been detected in rivers, lakes, and streams in 29 states, at levels that can result in detrimental effects to keystone aquatic organisms and the entire aquatic food web.²⁶

 Runoff from synthetic chemical fertilizers pollutes streams and lakes and causes algae blooms, depleted oxygen and damage to aquatic life.

THE REGISTRATION SYSTEM & PESTICIDE REGULATION

- The health data assessed by the U.S. Environmental Protection Agency (EPA) for the registration of pesticides comes from the manufacturer of the pesticide. EPA is not obligated under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA) to review peer-reviewed scientific literature.
- EPA often registers pesticides through a program called "conditional registration." In these cases, the agency permits a pesticide to go to market without all of its required data on health and environmental impacts because the agency assumes that no harm will result as it waits for this data. The U.S. Government Accountability Office (GAO) has criticized the agency for this process, noting that EPA "does not have a reliable system...to track key information related to conditional registrations."²⁷

- EPA's evaluation of endocrine (hormone) disrupting pesticides is years behind schedule and has been criticized for using outdated methods. The National Academy of Sciences has urged the agency to alter its approach to adequately address the low dose impacts of these chemicals.²⁸
- EPA only tests the active ingredient in pesticide formulations. Despite the fact that a pesticide product can contain multiple ingredients, the agency does not look at synergistic effects.
 Science shows that combinations of active ingredients can increase or decrease the toxicity of a product, but this impact is simply not evaluated by the agency.²⁹
- Most states have preemption laws that prohibit localities from passing ordinances that restrict pesticides on private property more stringently than the state policy.³⁰

The National Academy of Sciences has urged the agency to...address the low dose impacts of these chemicals.

"INERT" INGREDIENTS

- Pesticide products are made of an active ingredient and several "inert," or other, ingredients. "Inert" ingredients are not chemically, biologically, nor toxicologically inert. "Inerts" are not disclosed to the public due to their status as "trade secrets."
- Active ingredients usually comprise only 5% of the actual product; the other ingredients make up the majority of a given pesticide product or formulation.³¹
- "Inert" ingredients can be more toxic to humans than the active ingredient. Polyethoxylated tallow amine, or POEA, often found in Roundup formulations with glyphosate, is an example of an "inert" ingredient linked to damage to embryonic, placental and umbilical cord cells.³²
- After a 2006 proposal by EPA to disclose "inert" ingredients allowed in pesticide formulations, the agency retracted its original intent and released information about only 72 of the 371 "inerts" allowed in pesticide product formulations. EPA further indicated the disclosed "inerts" were no longer in use.³³

ENDNOTES

- U.S. Environmental Protection Agency (EPA). 2017. Pesticides Industry Sales and Usage: 2008-2012 Market Estimates. https://www.epa.gov/sites/production/ files/2017-01/documents/pesticidesindustry-sales-usage-2016 0.pdf.
- 2 Ibid.
- 3 National Research Council. 1980. Urban Pest Management. National Academy of Sciences; Abrams, R., Attorney General of New York. 1991. "Toxic Fairways: Risking Groundwater Contamination from Pesticides on Long Island Golf Courses," Environmental Protection Bureau; Pimentel, D, et al. 1991. "Environmental and Economic Impacts of Reducing U.S. Agricultural Pesticide Use," Handbook of Pest Management in Agriculture, 2nd ed. CRC Press, Florida, p.679.
- 4 United States Census Bureau. 2016. 2012 Economic Census- Landscaping Services. https://factfinder.census.gov/faces/ tableservices/jsf/pages/productview. xhtml?src=bkmk.
- 5 Ibid.

- 5 The National Gardening Association and Organic Gardening Magazine. 2004 July. Environmental Lawn and Garden Survey.
- 7 Beyond Pesticides Factsheet. 2015. Health Effects of 30 Commonly Used Lawn Pesticides. http://www.beyondpesticides. org/lawn/factsheets/30health.pdf.
- 6 EPA. 2003. Tackling a Suspected Hazard of Aging. http://www.epa.gov/ord/archives/2003/september/htm/article1. htm (accessed 3/4/05); U.S. EPA. 2002 Oct 31. "EPA Announces New Aging Initiative to Protect Older Persons From Environmental Health Threats." American Academy of Pediatrics. 2012. Pesticide Exposure in Children. Pediatrics. http://pediatrics.aappublications.org/content/130/6/e1757.
- 9 Rudel, Ruthann, et al. 2003. "Phthalates, Alkylphenols, Pesticides, Polybrominated Diphenyl Ethers, and Other Endocrine-Disrupting Compounds in Indoor Air and Dust," Environmental Science and Technology 37(20): 4543–4553; Nishioka, M., et al. 2001. "Distribution of 2,4-D in Air and on Surfaces Inside Residences After Lawn Applications: Comparing Exposure

- Estimates from Various Media for Young Children," Environmental Health Perspectives 109(11); Trunnelle, KJ et al. 2014. Urinary Pyrethroid and Chlorpyrifos Metabolite Concentrations in Northern California Families and Their Relationship to Indoor Residential Insecticide Levels, Part of the Study of Use of Products and Exposure Related Behavior (SUPERB). http://pubs.acs.org/doi/abs/10.1021/es403661a.
- 10 EPA. 2015. Pesticides and their Impact on Children: Key Facts and Talking Points. https://www.epa.gov/sites/production/ files/2015-12/documents/pest-impacthsstaff.pdf.
- 11 National Institutes of Health. 2008.
 President's Cancer Panel-Reports. Reducing
 Environmental Cancer Risk: What We
 Can Do Now. https://deainfo.nci.nih.gov/advisory/pcp/annualReports.
- 12 Green KR, Peters S, Bailey HD. 2013. Exposure to pesticides and the risk of childhood brain tumors. Cancer Causes and Control. Jul;24(7):1269–78 https:// www.ncbi.nlm.nih.gov/ pubmed/23558445.

- 13 Greenlee, A. et al. 2004. "Low-Dose Agrochemicals and Lawn-Care Pesticides Induce Developmental Toxicity in Murine Preimplantation Embryos," Environ Health Perspect 112(6): 703–709; Cavieres, M., et al. 2002. "Developmental toxicity of a commercial herbicide mixture in mice: Effects on embryo implantation and litter size." Environ Health Perspect 110: 1081–1085.
- 14 Salam, M.T., et al. 2004. "Early Life Environmental Risk Factors for Asthma: Findings from the Children's Health Study," Environ Health Perspectives 112(6): 760.
- 15 Wagner-Schuman, M, Richardson, J, Auinger, P et al. 2015. Association of pyrethroid pesticide exposure with attentiondeficit/hyperactivity disorder in a nationally representative sample of U.S. children. Environmental Health. 14:44.
- 16 Centers for Disease Control and Prevention. 2003 Jan. Second National Report on Human Exposure to Environmental Chemicals; Pohl, HR., et al. 2000. "Breastfeeding exposure of infants to selected pesticides," Toxicol Ind Health 16: 65-77; Sturtz, N., et al. 2000. "Detection of 2,4-Dichlorophenoxyacetic acid residues in neonates breast-fed by 2,4-D exposed dams," Neurotoxicology 21(1-2): 147-54; Houlihan, J., et al. 2005. Body Burden, The Pollution in Newborns. Environmental Working Group, Washington, D.C.
- 17 Takashima-Uebelhoer, BB et al. 2012. Household chemical exposures and the risk of canine malignant lymphoma, a model for human non-Hodgkin's lymphoma. Environmental Research. 112 (171–176).
- 18 Beyond Pesticides Factsheet. 2015. Environmental Effects of 30 Commonly Used Lawn Pesticides. http://www. beyondpesticides.org/lawn/factsheets/ 30enviro.pdf.
- 19 Defenders of Wildlife. The Dangers of Pesticides to Wildlife [white paper]. 2005 April. www.pesticidefreelawns.org/resources.

- 20 Restmeyer, S.J. 2003. Ecological Pest Management: Embracing the Organic Approach to Landscape Management. Pesticides and You 23(1): 11–12. Beyond Pesticides, Washington, D.C.
- 21 Toher et al. 2014. No Longer a Big Mystery. Pesticides and You. (34)(1) 9-12. Beyond Pesticides, Washington, D.C. http://www.beyondpesticides.org/assets/media/documents/pollinators/ nolongeraBIGmystery.pdf.
- 22 Pesticides Factsheet. 2005. Environmental Effects of 30 Commonly Used Lawn Pesticides. http://www.beyondpesticides.org/lawn/factsheets/30enviro.pdf.
- 23 Battaglin et al. 2014. Glyphosate and its Degradation Product AMPA Occur Frequently and Widely in U.S. Soils, Surface Water, Groundwater, and Precipitation. Journal of the American Water Resources Association. (50)(2) 275-290.
- 24 Ryberg, K.R., Vecchia, A.V., Martin, J.D., and Gilliom, R.J., 2010, Trends in pesticide concentrations in urban streams in the United States, 1992–2008: U.S. Geological Survey Scientific Investigations Report 2010–5139, 101; DeSimone, Leslie. 2009. United States Geological Survey. "Quality of Water from Domestic Wells in Principal Aquifers of the United States, 1991–2004." http://pubs.usgs.gov/sir/2008/5227/includes/sir2008-5227.pdf.
- 25 Ontario Ministry of the Environment. 2011. Pesticide Concentrations in Ontario's Urban Streams One Year After the Cosmetic Pesticides Ban. http://www.landscapeontario. com/attach/1295274330.MOE_Update_-Aaron Todd.pdf.
- 26 Harriott, Nichelle and Shistar, Terry. 2017. Poisoned Waterways: The same pesticide that is killing bees is destroying life in the nation's streams, rivers, and lakes. Pesticides and You. (40)(1). 9–18. Beyond Pesticides, Washington, D.C.
- 27 U.S. Government Accountability Office. 2013. EPA Should Take Steps to Improve Its Oversight of Conditional Registrations. GAO-13-145. http://www.gao.gov/ products/GAO-13-145.

- 28 Colborn, Theo. 2009. EPA's new pesticide testing is outdated, crude. Environmental Health News. http://www.environmental healthnews.org/ehs/editorial/epa2019s-new-pesticide-testing-is-outdated-crude; National Academies of Sciences, Engineering, and Medicine. 2017. Application of Systemic Review Methods in an Overall Strategy for Evaluating Low-Dose Toxicity from Endocrine Active Chemicals. https://www.nap.edu/catalog/24758/application-of-systematic-review-methods-in-an-overall-strategy-for-evaluating-low-dose-toxicity-from-endocrine-active-chemicals.
- 29 Center for Biological Diversity. 2016. Toxic Concoctions: How the EPA Ignores the Dangers of Pesticide Cocktails. http://www. biologicaldiversity.org/campaigns/ pesticides_reduction/pdfs/Toxic_ concoctions.pdf.
- 30 Porter, Matt. 2013. State Preemption Law: The battle for control of democracy. Pesticides and You. (33)(3). Beyond Pesticides, Washington D.C.
- 31 Spitzer, E., Attorney General of NY. 2000. The Secret Ingredients in Pesticides: Reducing Risk. Abrams, R., 1991. Attorney General of NY. "The Secret Hazards of Pesticides: Inert Ingredients."
- 32 Benachour, N and Seralini, G.E. 2009. Glyphosate Formulations Induce Apoptosis and Necrosis in Human Umbilical, Embryonic, and Placental Cells. Chemical Research and Toxicology. 22 (1), 97–105. http://pubs.acs.org/doi/abs/10.1021/ tx800218n.
- 33 EPA. 2014. EPA Proposes to Remove 72
 Chemicals from Approved Pesticide Inert
 Ingredient List. https://yosemite.epa.gov/
 opa/admpress.nsf/0/3397554fa65588d6
 85257d7a0061a300?OpenDocument;
 Beyond Pesticides. 2014. Daily News Blog.
 Groups Call for Labeling of 300 Inert
 Ingredients as EPA Delists 72 Already
 Discontinued. http://beyondpesticides.org/
 dailynewsblog/2014/10/epa-delisting-of72-inert-ingredients-no-longer-usedcalled-inadequate-response-to-problem.

